

Technical Bulletin

Genes that fit *your* farm.

SeCan

Canada's Seed Partner

AAC Starbuck VB Canada Western Red Spring Wheat



Progress Through Research
Le progrès grâce à la recherche



Description:

AAC Starbuck VB is a very high yielding semi-dwarf CWRS with good lodging tolerance and is rated MR to FHB resistance. AAC Starbuck VB is also tolerant to the orange wheat blossom midge. AAC Starbuck VB is well-adapted to the wheat growing regions across the western Canadian Prairies.

Certified seed of AAC Starbuck VB will be sold as a varietal blend made up of 90% AAC Starbuck and 10% AAC Brandon. Blending with the midge susceptible variety AAC Brandon provides a refuge area for non-virulent midge to survive at low levels, thereby extending the useful life of the *Sm1* midge tolerant gene.

Parentage: Griffon/BW876//Carberry

Strengths:

- 114% of the grain yield of AC[®] Carberry
- Semi-dwarf with height similar to AC[®] Carberry
- Moderately resistant to FHB, low DON accumulation
- Moderately resistant to leaf rust, stripe rust and loose smut

Neutral Traits:

- Grain protein potential similar to AC[®] Carberry
- Maturity similar to AC[®] Carberry
- Slightly weaker straw strength compared to AC[®] Carberry
- Intermediate resistance to stem rust

Weakness:

- Susceptible to common bunt and leaf spot

Breeder:

Dr. Richard Cuthbert
Swift Current Research and Development Centre
Agriculture and Agri-Food Canada
Swift Current, SK

PBR 91 Applied For

2015-2017 Western Bread Wheat Registration Trials - Registration Data

Variety	Yield (% AC Carberry)	Maturity (days)	Lodging 1 = erect 9 = flat	Height (cm)	Test Weight (kg/hl)	Kernel Weight (mg/kernel)	Grain Protein (%)	FHB Resistance Rating
AC [®] Carberry	100	100.2	1.8	82	80.3	34.6	14.6	MR
AC [®] Unity VB	104	96.9	3.4	93	80.4	33.0	14.3	I
AAC Viewfield	109	100.4	1.7	77	81.1	32.6	14.6	I
Glenn	98	100	2.1	88	82.7	33.5	14.6	I
AAC Starbuck VB	114	100.8	2.1	82	81.1	35.1	14.6	MR

VB = varietal blend R=Resistant; MR=Moderately Resistant; I=Intermediate; MS=Moderately Susceptible; S=Susceptible

'AC' is an official mark used under license from Agriculture & Agri-Food Canada

For more information, call 1-800-665-7333 or visit www.secan.com

2020 Seed Manitoba - Wheat Comparison

Variety	Site Years Tested	Yield bu/ac	Protein %	Maturity +/- 99 days	Height +/- 81cm	Spike Awned	Resistance to:								
							Lodging	Sprouting	Loose Smut	Common Bunt	Leaf Spot	Stem Rust	Leaf Rust	Stripe Rust	FHB
AAC Alida VB*	33	68	14.6	+2	+6	Y	VG	VG	R	I	MS	R	R	MR	MR
AAC Brandon	61	70	14.3	+2	0	Y	VG	P	MR	S	I	R	R	MR	MR
AAC Tisdale	32	68	15.2	+1	+8	Y	G	F	MR	MR	MS	R	R	S	MR
AAC Warman VB*	33	68	14.4	0	+12	Y	G	---	MR	S	I	R	R	MS	MR
AAC Wheatland VB*	20	74	14.3	1	0	Y	VG	---	R	MR	S	R	R	I	I
AC® Carberry	124	67	14.6	2	0	Y	VG	F	MR	R	MS	MR	R	MR	MR
AAC Starbuck VB*	20	73	14.4	1	1	Y	G	---	MR	S	S	I	MR	MR	MR

*Varietal Blend F=Fair; G=Good; VG=Very Good R=Resistant; MR=Moderately Resistant; I=Intermediate; MS=Moderately Susceptible; S=Susceptible

2020 Varieties of Grain Crops for Saskatchewan - Wheat Comparison

Variety	Years Tested	Yield as % of Carberry			Protein	Resistance to:									Relative Maturity (days)	Head Awedness	Seed Weight (mg)	Test Weight (kg/hl)	Height (cm)
		Area 1 & 2	Area 3 & 4	Irrigation		Lodging	Sprouting	Stem Rust	Leaf Rust	Stripe Rust	Loose Smut	Bunt	Leaf Spot	FHB					
AC® Carberry	6	100	100	100	14.5	VG	F	MR	R	MR	MR	R	MS	MR	102	Y	35.8	80.3	83
AAC Alida VB*	3	106	108	---	-0.2	VG	VG	R	R	MR	R	I	MS	MR	0	Y	+1.5	+0.3	+7
AAC Brandon	5	106	106	---	-0.4	G	P	R	R	MR	MR	S	I	MR	0	Y	+0.1	0.0	-1
AAC Jatharia VB*	5	108	114	---	-0.2	F	G	I	R	I	S	MS	I	I	0	Y	+0.7	+0.9	+15
AAC Warman VB*	2	102	107	---	-0.3	F	F	R	R	MS	MR	S	I	MR	-1	Y	-1.6	+0.2	+13
AAC Wheatland VB*	2	111	113	---	-0.4	VG	G	R	R	I	R	MR	S	I	0	Y	-0.1	0.0	+2
AC® Shaw VB*	6	112	114	103	-0.7	F	G	R	MR	I	S	MR	MS	MS	-1	N	+0.5	-0.5	+18
AAC Starbuck VB*	2	113	117	---	-0.5	G	F	I	MR	MR	MR	S	S	MR	0	Y	+0.2	+0.5	+3

*Varietal Blend G=Good; VG=Very Good; F=Fair; P=Poor; VP=Very Poor

2020 Alberta Seed Guide - CWRS Wheat Comparison

Variety	Overall Yield		Test Yield Category			Maturity Rating (Days +/- Carberry)	Protein %	Test Weight (lb/bu)	Kernel Weight g/1000	Height (cm)	Resistance to:		Disease Resistance:						
	All Sites	Station years of testing	Low < 55 bu/ac	Med 55 - 80 bu/ac	High >80 bu/ac						Ldg.	Sprout	Loose Smut	Bunt	Stripe Rust	Leaf Spot	FHB		
			Yield as % of AC® Carberry																
AC® Carberry bu/ac	70		45	66	95														
AC® Carberry	100	126	100	100	100	104	13.9	63	40	80	VG	F	MR	R	MR	MS	MR		
AAC Alida VB*	101	45	98	103	99	0	0	63	41	88	G	VG	R	I	MR	MS	MR		
AAC Brandon	105	35	XX	103	107	0	-0.3	63	41	84	G	P	MR	S	MR	I	MR		
AAC Tisdale	101	45	100	101	102	-1	+0.4	63	42	88	F	F	MR	MR	S	MS	MR		
AAC Wheatland VB*	107	31	XX	105	109	0	-0.5	63	41	85	VG	G	R	MR	I	S	I		
AC® Shaw VB*	105	53	104	105	107	-1	-0.9	63	37	92	G	G	S	S	MR	I	MS		
AAC Starbuck VB*	106	31	XX	106	108	0	-0.4	62	41	86	G	F	MR	S	MR	S	MR		

*Varietal Blend VG = Very Good; G = Good; F = Fair; P = Poor; VP = Very Poor; Disease Ratings: R=Resistant; MR=Moderately Resistant; I=Intermediate; MS=Moderately Susceptible; S=Susceptible

For more information, call 1-800-665-7333 or visit www.secan.com